
Social Support and Oxytocin Interact to Suppress Cortisol and Subjective Responses to Psychosocial Stress

Markus Heinrichs, Thomas Baumgartner, Clemens Kirschbaum, and Ulrike Ehlert

Background: *The presence of social support has been associated with decreased stress responsiveness. Recent animal studies suggest that the neuropeptide oxytocin is implicated both in prosocial behavior and in the central nervous control of neuroendocrine responses to stress. This study was designed to determine the effects of social support and oxytocin on cortisol, mood, and anxiety responses to psychosocial stress in humans.*

Methods: *In a placebo-controlled, double-blind study, 37 healthy men were exposed to the Trier Social Stress Test. All participants were randomly assigned to receive intranasal oxytocin (24 IU) or placebo 50 min before stress, and either social support from their best friend during the preparation period or no social support.*

Results: *Salivary free cortisol levels were suppressed by social support in response to stress. Comparisons of pre- and poststress anxiety levels revealed an anxiolytic effect of oxytocin. More importantly, the combination of oxytocin and social support exhibited the lowest cortisol concentrations as well as increased calmness and decreased anxiety during stress.*

Conclusions: *Oxytocin seems to enhance the buffering effect of social support on stress responsiveness. These results concur with data from animal research suggesting an important role of oxytocin as an underlying biological mechanism for stress-protective effects of positive social interactions.* Biol Psychiatry 2003;54:1389–1398 © 2003 Society of Biological Psychiatry

Key Words: Social support, oxytocin, stress, cortisol, anxiety, intranasal

Introduction

There is substantial evidence indicating that psychosocial risk factors (e.g., lack of social contact, death of a spouse) contribute to a wide spectrum of somatic, psychosomatic, and psychiatric disorders with major public health significance, such as depression or coronary heart disease (Bruce 2002; Ehlert and Straub 1998; Kraemer et al 2001; Rozanski et al 1999; Ruberman et al 1984; Smith and Ruiz 2002). Positive social interactions have in turn been shown to exert powerful beneficial effects on health outcomes and longevity. Social support is the most intensively investigated social factor in humans (Broadhead et al 1983; House et al 1988; Knox and Uvnas-Moberg 1998; MacMahon and Lip 2002; Paykel 2001; Schwarzer and Leppin 1991; Seeman 2000; Seeman and McEwen 1996; Uchino et al 1996; Veiel and Baumann 1992). A growing body of research in clinical populations has provided support for the hypothesis that higher reported levels of social support are associated with positive effects on various diseases, such as cardiovascular reactivity and blood pressure (Evans and Steptoe 2001; Gallo et al 2000; Spitzer et al 1992; Steptoe 2000; Uchino et al 1999; Uno et al 2002), depression (Hays et al 2001; Sayal et al 2002), and schizophrenia (Buchanan 1995; Erickson et al 1998; Macdonald et al 1998). In addition, physiologic stress reactivity in experimental studies has also been shown to be responsive to social support. The availability of social support has been associated with attenuated free cortisol concentrations in saliva (Kirschbaum et al 1995) and lower cardiovascular reactivity (Gerin et al 1992; Lepore et al 1993; Uchino and Garvey 1997) in response to acute laboratory stress.

Although there is considerable knowledge concerning the positive effects of social support on morbidity and mortality risks, to date, less attention has been given to the underlying physiologic mechanisms or pathways. More specifically, it is not clear how positive social interactions suppress stress-responsive physiologic systems or stimulate other internal regulatory systems involved in the attenuation of stress reactivity. A better understanding of these mechanisms would undoubtedly have important

From the Department of Clinical Psychology (MH, TB, UE), University of Zürich, Zürich, Switzerland; Center for Psychobiological and Psychosomatic Research (MH), University of Trier, Trier, Germany; and Institute of Experimental Psychology II (CK), University of Düsseldorf, Düsseldorf, Germany.

Address reprint requests to Markus Heinrichs, Ph.D., University of Zürich, Department of Clinical Psychology, Zürichbergstr. 43, CH-8044 Zürich, Switzerland.

Received February 7, 2003; revised April 16, 2003; accepted April 17, 2003.

clinical implications for differential diagnosis and specific psychological and pharmacologic treatments for numerous disorders.

In animals, the neuropeptide oxytocin has been shown to exert behavioral and physiologic stress-attenuating and anxiolytic effects and, in addition, might prove to promote positive social interaction (Carter 1998; Carter and Altemus 1997; Legros 2001; McCarthy and Altemus 1997; Pedersen 1997; Pedersen and Boccia 2002; Uvnas-Moberg 1998). Oxytocin is synthesized in magnocellular neurons of the paraventricular and supraoptic nuclei of the hypothalamus (Swaab et al 1975; Vandesande and Dierickx 1975). It is processed from its precursor form, together with the carrier protein, along the axonal projection to the posterior pituitary, from which the peptide is secreted into the systemic circulation (Brownstein et al 1980). In addition, oxytocin is widely distributed throughout the central nervous system from smaller parvocellular neurons, influencing many neurobehavioral functions (de Wied et al 1993; McCarthy and Altemus 1997). In experimental studies in animals, oxytocin has been stimulated with both exogenous (e.g., intracerebroventricular injection) and endogenous (e.g., suckling stimulus during lactation) methods of stimulation. Aside from the most well-known peripheral role of oxytocin in parturition and lactation, intracerebral oxytocin inhibits the stress-induced activity of the hypothalamic–pituitary–adrenal (HPA) axis responsiveness, suggesting an inhibitory influence of oxytocin on stress-responsive neurohormonal systems (Neumann et al 2000a, 2000b; Uvnas-Moberg et al 1994, 1999; Windle et al 1997b). Moreover, the neuropeptide has been shown to function as a central regulator in social attachment and in related prosocial behaviors (Carter 1998; Insel 1997; Insel and Young 2001; Pedersen 1997).

In contrast to the very active investigation in recent years into the effects of oxytocin on physiologic and behavioral stress and anxiety responses in animals, research in humans remains relatively limited; however, initial studies suggest similar stress-reducing effects of the neuropeptide in humans. Similar to the aforementioned studies in lactating rodents, the lactation paradigm for endogenous stimulation has also been used in human studies. In lactating women, a suppression of endocrine stress responses has been observed if breast-feeding starts 30–60 min before stress exposure, depending on the kind of stressor (Altemus et al 1995, 2001; Heinrichs et al 2001, 2002). Within this context, however, it should be noted that there are a variety of confounding factors, in particular the release of other hormones, that are difficult to control for in endogenous stimulation paradigms. Thus, the specific effects of central oxytocin as an underlying biological mechanism for the reduction of stress and anxiety and for positive social interactions in humans are yet to be

determined. Alternatively, studies on central effects of oxytocin and vasopressin in healthy humans and patients have been carried out with exogenous stimulation with intranasal administration (Born et al 1998; Bruins et al 1992; Heinrichs 2000; Pitman et al 1993). Neuropeptides have recently been shown to enter the cerebrospinal fluid directly after intranasal administration (Born et al 2002). To date, there have been no human studies that directly address effects of oxytocin administration and social support on stress and anxiety. With these considerations in mind, we set out to study the effects of oxytocin and social support on mood, anxiety, and neuroendocrine responses to psychosocial stress (public speaking and mental arithmetic in front of an audience) in healthy men.

Methods and Materials

Participants

Thirty-seven healthy men (mean \pm SD age, 23.8 ± 3 years) were recruited by advertisements posted at the university and were paid for participation. The study was conducted at the University of Trier, Germany. All subjects underwent a physical evaluation to screen out chronic diseases, mental disorders, medication, smoking, and drug or alcohol abuse. Participants abstained from food and drink (other than water) for 2 hours before the experiment, and from exercise, caffeine, and alcohol during the 24 hours before the session. Three of the original 40 subjects were excluded: one who scored at 65 (the 100th percentile) in the State-Trait-Anxiety Inventory (Laux et al 1981) and two who were found to have taken medication because of acute seasonal allergic rhinitis. The study was carried out in accordance with the Declaration of Helsinki principles and approved by the institutional ethics committee. All subjects gave written, informed consent and were informed of their right to discontinue participation at any time.

Social Support Provision

Two weeks before the experiment, subjects were randomly instructed in writing either to bring their best friend (male or female) along with them to the experimental session (“social support”) or to come alone (“no social support”). Support providers were told that their task during the experiment was to be as helpful as possible during the 10-min preparation for the speech task (see Stress Procedure) and offer both instrumental and emotional support. More specifically, they were told that they would know best what kind of supportive behaviors would fit the individual coping preferences of the subject. Support providers were present only during preparation of the stressor.

Oxytocin Administration

Participants received a single dose of 24 IU oxytocin (Syntocinon-Spray, Novartis, Basel, Switzerland) intranasally or placebo. Intranasal administration took place immediately after the introduction at the beginning of the experiment (50 min before stress exposure); a dose of 12 IU was sprayed into each nostril (three

puffs per nostril, each with 4 IU oxytocin). The methodology of the administration used has been described elsewhere (Heinrichs 2000). Intranasal oxytocin is widely prescribed in lactating women and is well tolerated. Several studies have been conducted in humans with doses between 20 and 60 IU, and no adverse side effects have been reported (e.g., Bruins et al 1992; Fehm-Wolfsdorf et al 1988; Pitman et al 1993).

Stress Procedure

All experimental sessions commenced between 2:00 PM and 4:00 PM to control for diurnal changes in cortisol secretion and lasted for approximately 2 hours. Based on random assignment, participants either brought their best friend along with them or came alone to the session. The Trier Social Stress Test (TSST; Kirschbaum et al 1993), which primarily consists of a public speaking task and mental arithmetic performed in front of an audience, was used for stress induction. This standardized psychosocial stress induction technique enables a naturalistic exposure to a psychosocially stressful situation. At the time of recruitment into the study, participants were informed that the experiment consisted of a public speaking task during which saliva samples would be collected for the measurement of the stress hormone cortisol. During the introduction to the TSST they were then told that they would be required to give a 5-min mock job interview to an unknown panel (consisting of one man and one woman) on personal suitability for a job and to enumerate their strengths and qualifications in an unstructured manner, followed by 5 min of mental arithmetic performed out loud. To increase task engagement, the job description was matched to each participant, taking into consideration his own individual goals and aspirations. The panel of evaluators were presented as experts in the evaluation of nonverbal behavior.

After being shown the TSST room containing the panel of evaluators and a conspicuous video camera, subjects were given 10 min for either solitary or socially supported preparation for this task (see Social Support Provision). After this, subjects re-entered the TSST room and were told to start the presentation. Any pause during the speech resulted in the participant being reminded of the remaining time. At the end of the interview, the subjects were instructed to serially subtract 13 from 2034 as quickly and accurately as possible, still in front of the panel. If the subject made a mistake, he was told to start over from the beginning. After 5 min of mental arithmetic, the interview was concluded.

Previous studies have indicated that this stress protocol reliably induces a significant activation of physiologic and psychological responses to stress in healthy subjects and patients (e.g., Domes et al 2002; Hellhammer et al 1997; Heinrichs et al 2001; Kirschbaum et al 1999; Young et al 2000). Subjects were confronted with the psychosocially stressful situation for a total of 10 min. At the end of the saliva sampling, participants were debriefed and given the name of a contact person should they have any further questions.

Salivary Cortisol Measures

Salivary cortisol is considered a reliable and valid measure of unbound ("free") cortisol concentrations in plasma (Vining and

McGinley 1987). A commercially available sampling device (Salivette; Sarstedt, Rommelsdorf, Germany) was used to obtain eight saliva samples from each participant over the course of the stress session. Saliva samples were collected before the preparation phase (baseline: -20 min relative to TSST) and immediately before stress (pretrial: -1 min). Six additional samples were collected during the 60-min poststress period (+1, +10, +20, +30, +45, and +60 min).

After being chewed for about 60 sec, devices were stored at -20°C until required for biochemical analysis. Before assaying for free cortisol, samples were thawed and spun at 3000 rpm for 5 min to obtain .5–1.0 mL clear saliva with low viscosity. The free cortisol concentration in saliva was analyzed with a time-resolved immunoassay with fluorescence detection, as described previously (Dressendörfer et al 1992). The limit of detection was .5 nmol/L for saliva cortisol. Inter- and intra-assay coefficients of variance were below 12% and 10%, respectively.

Psychological Measures

Depression and anxiety were assessed by the German versions of the Self-Rating Depression Scale (Zung 1965) and the State-Trait Anxiety Inventory (Laux et al 1981). To assess the general availability of social support, we used the German version of the Interpersonal Support Evaluation List (Laireiter 1996). This questionnaire measures the perceived availability of social support resources. We used a mood questionnaire that is especially suited for repeated measures within several minutes or hours (Steyer et al 1997). Twelve items are rated on a 5-point scale, ranking from 1 ("not at all") to 5 ("very strongly"). Factor analysis revealed three scales, termed elevated versus depressed mood, wakefulness versus sleepiness, and calmness versus restlessness. Self-reported affect was assessed by measuring mood and state anxiety before and after stress exposure.

Data Analyses

For each variable of interest, we used three-way analyses of variance (ANOVA) with repeated measurement (social support \times oxytocin \times time [repeated factors: eight for cortisol and two for anxiety and mood, respectively]) (Winer 1971). When the Mauchly test of sphericity indicated heterogeneity of covariance, we verified repeated-measures results with Greenhouse-Geisser corrections. To determine where significant differences from baseline hormone levels in groups were occurring, multiple single comparisons were conducted with an adjusted level of significance (*t* test with Bonferroni corrections). The areas under the individual response curves were calculated with the trapezoid formula, aggregating the eight saliva hormone levels during the stress protocol. Statistical tests were two-tailed and conducted at the .05 level of significance.

Results

Description of the Study Groups

As described in Methods and Materials, all participants were randomly assigned to receive intranasal oxytocin (24 IU) or placebo 50 min before stress, and either no social support or social support from their best friend during the

Table 1. Characteristics of Study Groups

	No Social Support and Placebo (n = 8)	Social Support and Placebo (n = 10)	No Social Support and Oxytocin (n = 8)	Social Support and Oxytocin (n = 11)
Age (years)	26.0 ± 1.0	23.3 ± .9	22.6 ± 1.0	23.5 ± .9
Body Mass Index (kg/m ²)	23.6 ± .7	22.4 ± .6	22.2 ± .7	22.3 ± .6
Social Support (ISEL score)	71.8 ± 4.5	77.0 ± 4.0	70.4 ± 4.5	75.4 ± 3.8

Data given as mean ± SEM. ISEL, Interpersonal Support Evaluation List.

10-min preparation period (placebo-controlled, double-blind, 2 × 2 factorial design). As shown in Table 1, subjects of the four study groups did not differ significantly with respect to age and body mass index. There were no differences in the perceived availability of general social support among groups. Trait anxiety (36.5 ± 1.0) and depression scores (40.6 ± 1.1) of all participants were within the normal range for the general population.

Cortisol Responses to Stress

In the overall analyses, three-way analysis of variance with repeated measurement revealed the expected significant salivary free cortisol response to stress in the total group [time effect: $F(2.4,80.5) = 37.40, p < .001$]. Figure 1 depicts the means and standard errors among groups for saliva cortisol at each time point and the area under the response curves. No significant differences in pretrial cortisol levels were observed among groups 1 min before stress. There was a significant attenuating effect of social support on cortisol [social support × time effect, $F(2.4,80.5) = 8.52, p < .001$] and a trend toward

attenuated cortisol levels by oxytocin [oxytocin × time effect, $F(2.4,80.5) = 2.45, p = .08$]. More importantly, a significant three-way interaction effect was obtained [social support × oxytocin × time effect: $F(2.4,80.5) = 4.54, p < .01$], with the lowest cortisol concentrations during stress in subjects who received both social support and oxytocin.

The mean absolute increase in salivary cortisol in response to stress was 15.1 nmol/L in subjects without social support and with placebo, 3.6 nmol/L in subjects with social support and placebo, 6.7 nmol/L in subjects without social support and with oxytocin, and 4.0 nmol/L in subjects with both social support and oxytocin. Subjects without social support and with placebo showed significantly higher cortisol levels at 1, 10, 20, 30, 45, and 60 min after cessation of stress compared with pretrial levels 1 min before stress (all $p < .05$ with Bonferroni correction), whereas subjects without social support and with oxytocin yielded significant cortisol increases at 1, 10, and 20 min after stress exposure (all $p < .05$ with Bonferroni correction). Notably, no significant cortisol increases after stress in comparison with pretrial concentrations occurred




Figure 1. Mean salivary free cortisol concentrations (± SEM) during psychosocial stress exposure (Trier Social Stress Test). Participants were randomly assigned to receive intranasal oxytocin (24 IU) or placebo and either no social support or social support from their best friend before stress. The shaded area indicates the period of a public speaking task followed by mental arithmetic in front of a panel of evaluators. The areas under the individual response curves aggregate the eight saliva hormone levels during the stress protocol. Significant interaction effect on cortisol was obtained (social support × time effect, $p < .001$; social support × oxytocin × time effect, $p < .01$).




Figure 2. Mean (with SEM bars) levels of calmness as a function of study group and time of measurement (before and after stress). Significant three-way interaction effect on calmness was obtained (social support \times oxytocin \times time effect, $p < .05$).

in participants with social support and placebo or in participants with social support and oxytocin.

Mood and Anxiety Responses to Stress

Mood and anxiety were assessed before and after the psychosocial stress test. Analysis of variance with repeated measurement revealed a significant three-way interaction effect on calmness [social support \times oxytocin \times time effect: $F(1,33) = 4.14$, $p = .05$] (Figure 2). No significant differences in calmness before stress induction were observed among groups. As expected, subjects without social support and with placebo exhibited a decrease in calmness during stress (before: 13.3 ± 1.4 ; after: 11.3 ± 1.7). In contrast, participants who received either social support (before: $9.8 \pm .9$; after: 12.8 ± 1.2) or oxytocin (before: $12.9 \pm .9$; after: $14.4 \pm .7$) or both social support and oxytocin (before: 12.6 ± 1.4 ; after: $13.9 \pm .9$) showed increasing calmness during the stress procedure (Figure 2). No changes in the mood and wakefulness scales were observed.

A trend toward a three-way interaction on state anxiety was obtained [social support \times oxytocin \times time effect: $F(1,33) = 4.05$, $p = .052$]. Again, there were no significant differences in anxiety scores among groups before stress. The group without social support and with placebo showed the expected increase in anxiety (before: 39.0 ± 2.9 ; after: 41.8 ± 4.3), whereas participants with either

social support (before: 47.4 ± 3.0 ; after: 43.0 ± 2.8) or oxytocin (before: 42.9 ± 2.6 ; after: 38.1 ± 2.3) or both factors (before: 43.0 ± 3.2 ; after: 41.5 ± 1.9) showed decreasing anxiety during stress. Importantly, comparisons of pre- and poststress anxiety levels revealed a significant anxiolytic effect of intranasal oxytocin administration within the group that received oxytocin but not social support [$t(7) = 2.68$, $p < .05$].

Discussion

This is the first study on the effects of social support and oxytocin on endocrine, mood, and anxiety responses to psychosocial stress in humans. Participants who received both protective factors of social support and oxytocin exhibited the lowest cortisol concentrations during stress exposure, whereas subjects who received no social support and placebo demonstrated the highest cortisol response (Figure 1). Most notably, we found corresponding results in psychological measures, indicating that subjects without social support and with placebo showed the expected decrease in calmness and an increase in anxiety during stress. In contrast, participants who received either social support or oxytocin or both protective factors showed increasing calmness and decreasing anxiety scores during the stress procedure (Figure 2). Moreover, pre- and post-stress comparisons of anxiety showed an anxiolytic effect of oxytocin administration. The present data build on prior research by demonstrating effects of social support both at the physiologic and the psychological level of analysis and, in addition, by experimentally varying a possible underlying endocrine mechanism, the neuropeptide oxytocin. It might be concluded that the interaction effect of oxytocin and social support on both endocrine and subjective measures accounts for a central nervous mechanism in humans.

In the present study, we obtained further evidence of blunted physiologic stress responsiveness by social support in men, replicating previous reports on attenuated salivary cortisol levels (Kirschbaum et al 1995) and lower cardiovascular reactivity (Gerin et al 1992; Lepore et al 1993; Uchino and Garvey 1997) in response to a psychosocial stressor. Once again, it is acute available instrumental and emotional support offered by another person that is associated with a suppression of stress-responsive physiologic systems (in this case, free cortisol in saliva; Figure 1). Socially supported subjects might appraise the anticipated acute stressor as less threatening and more controllable than unsupported subjects (Kirschbaum et al 1995). These results from laboratory research are in accordance with clinical studies indicating numerous health benefits of social support (Buchanan 1995; Erickson et al 1998; Evans and Steptoe 2001; Gallo et al 2000; Hays et al 2001;

Macdonald et al 1998; Sayal et al 2002; Steptoe 2000; Uchino et al 1999; Uno et al 2002). This consistent finding speaks for the unique nature of positive social interaction that might modulate the activity of the HPA axis and the autonomic nervous system.

In animal research, it has been suggested that the central oxytocinergic system mediates this buffering effect on stress (Carter 1998; Uvnas-Moberg 1998). Several studies have been published in which the effects of oxytocin on stress and anxiety responses are investigated. Oxytocin is released peripherally and within the brain in response to both physical and psychological stress and fearful situations (Neumann et al 2000a, 2000b). The stimulated release of the neuropeptide in rodents has been shown to ameliorate the symptoms associated with anxiety and stress (Carter and Altemus 1997). In addition, central administration of oxytocin in male rats has been shown to have similar effects to benzodiazepines (Neumann et al 2000a, 2000b; Uvnas-Moberg et al 1994, 1999; Windle et al 1997a). Lactating animals provide an additional paradigm for examining the consequences of peripheral and central release of oxytocin. During lactation, the suckling stimulus by the newborn increases both oxytocin and prolactin release and decreases basal plasma levels of adrenocorticotropin (ACTH) and cortisol, suggesting an inhibitory influence of both peptides on stress-responsive neurohormonal systems (Amico et al 1994; Chiodera et al 1991; Heinrichs et al 2001, 2002; Nissen et al 1996; Russell et al 2001). Numerous observations in the lactating rat have shown a neuroendocrine hyporesponsiveness to physical and psychological situations that induce anxiety and stress, including attenuated secretion of ACTH (Lightman and Young 1989; Neumann et al 1998; Walker et al 1992, 1995; Windle et al 1997b), corticosterone (Neumann et al 1998; Walker et al 1992; Windle et al 1997b), and catecholamines (Higuchi et al 1989). In addition, data from different species and paradigms implicate oxytocin in social attachment and in related prosocial behaviors (Carter 1998; Insel 1997; Insel and Young 2001; Pedersen 1997). Together, the inhibitory effect of intracerebral oxytocin on stress-induced activity of HPA axis responsiveness and prosocial behavior in both male and female rodents points to its key role in behavior and stress management.

In accordance with these findings in animals, we recently demonstrated attenuated pituitary–adrenal reactivity to psychosocial stress after endogenous stimulation of oxytocin after breast-feeding in postpartum lactating women (Heinrichs et al 2001). In addition, lactating women had reduced plasma ACTH, cortisol, and glucose responses to physical stress in comparison with postpartum nonlactating women (Altemus et al 1995). Breast-feeding mothers with increased plasma oxytocin in re-

sponse to a speech stressor that immediately followed baby holding had lower blood pressure than did mothers with a decrease in oxytocin after stress (Light et al 2000). Moreover, nonpostpartum healthy women who showed increased plasma oxytocin levels in response to positive emotion and massage as well as maintained oxytocin levels during negative emotion were less likely to report interpersonal problems associated with intrusiveness (Turner et al 1999). Finally, studies on psychiatric disorders show decreased oxytocin concentrations, for example in plasma in patients with autism (Green et al 2001; Modahl et al 1998), in plasma in high-scoring pain and depression in fibromyalgia syndrome patients (Anderberg and Uvnas-Moberg 2000), and in cerebrospinal fluid in patients with anorexia nervosa and bulimia nervosa (Demitrack et al 1990). Interestingly, cerebrospinal fluid oxytocin was normal in subjects after recovery from bulimia nervosa and anorexia nervosa (Frank et al 2000). In contrast, increased concentrations of oxytocin have been observed in cerebrospinal fluid of patients with Prader-Willi syndrome and obsessive-compulsive disorder (Leckman et al 1994; Martin et al 1998), whereas other studies could not find differences between patients with obsessive-compulsive disorder and healthy control subjects (Altemus et al 1999). Although it is difficult to draw conclusions about oxytocin functioning in specific brain regions from peripheral or cerebrospinal fluid measurement, the data do suggest that dysfunctions of oxytocin metabolism might be related to several psychiatric disorders.

Despite the stress-buffering effect of endogenous stimulation of peripheral oxytocin concentrations, it is difficult to control for the potential impact of confounding factors, especially other hormones, such as prolactin or opioid peptides. By using double-blind, placebo-controlled intranasal administration in our study, we circumvented the difficulties of unspecific stimulation paradigms. Several pathways to the brain have been demonstrated to pass the blood–brain barrier in neuropharmacologic research, and there is convincing evidence that peptides gain access to the brain after intranasal administration, bypassing the bloodstream (e.g., extracellular pathway, intraneuronal transport; Born et al 1998, 2002; Heinrichs 2000). Indeed, studies on intranasal oxytocin administration have been reported showing amnesic effects on memory performance (Bruins et al 1992) and inhibited physiologic responding in posttraumatic stress disorder (Pitman et al 1993), although studies on obsessive-compulsive disorder have revealed no effects of intranasal oxytocin (den Boer and Westenberg 1992; Epperson et al 1996). Overall, a great deal remains to be specified regarding adequate measurement of centrally active oxytocin in humans. Controversy exists, for example, as to the correlation between oxytocin

concentrations in blood and brain (Amico et al 1990; Engelmann et al 2000; Neumann et al 1993; Wotjak et al 1998). Human studies that include measurement of oxytocin concentrations in cerebrospinal fluid during social interaction and stress exposure are needed to further evaluate the central nervous role of the neuropeptide.

We have hypothesized that social support might attenuate endocrine and subjective responses to psychosocial stress. As known from animal research, this stress-protective effect of social support might be mediated through increased oxytocin concentrations, which were experimentally varied by placebo-controlled, double-blind intranasal administration. Indeed, we found that social support alone attenuated cortisol stress responses to a greater extent than did oxytocin alone. On the other hand, intranasal oxytocin had an anxiolytic effect with regard to pre- and poststress comparisons of anxiety levels; however, both social support and oxytocin together showed the highest stress-protective and anxiolytic effect. In sum, the effectiveness of social support might be connected with a sufficient central nervous availability of oxytocin. We therefore suggest that the peptide hormone might prove to be important in the stress-protective effect of positive social interaction, and, in turn, a dysregulated oxytocin metabolism might be associated with clinical disorders of psychosocial relevance. Future studies should explore whether patients with psychiatric disorders that are associated with social deficits (e.g., social phobia, autism) or patients with stress-related disorders could have characteristic dysfunctions in oxytocin metabolism. Prospective longitudinal data might help to model this potential interaction linking social behavior, oxytocin, and stress responsiveness. Furthermore, more direct investigations of central nervous system oxytocin function, including neuroimaging (e.g., radioactive labeling of the neuropeptide in positron emission tomography), are warranted. Finally, the neuropeptide might be a novel target for neuropsychopharmacologic research.

We thank the study participants and acknowledge the technical contributions and assistance of (in alphabetical order) Jutta Bennecke, Uta Berger, Begga Beyersmann, Ulrike Bucerius, Bernhild Doll, Anuschka Enzler, Annemarie Fritzen, Sarah Mannion, Gunther Meinschmidt, Rita Praprotnik, Ingrid Rummel-Frühau, Katrin Ruwisch, Silke Welter, Cornelia Zenglein, and Britta Zimmermann. The study was conducted at the Center for Psychobiological and Psychosomatic Research, University of Trier, Germany. Our grateful thanks to Professor Dirk H. Hellhammer for his support. We thank Dr. Dominique J. de Quervain for reviewing an early version of the manuscript.

References

- Altemus M, Deuster PA, Galliven E, Carter CS, Gold PW (1995): Suppression of hypothalamic-pituitary-adrenal axis

- responses to stress in lactating women. *J Clin Endocrinol Metab* 80:2954–2959.
- Altemus M, Jacobson KR, Debellis M, Kling M, Pigott T, Murphy DL, Gold PW (1999): Normal CSF oxytocin and NPY levels in OCD. *Biol Psychiatry* 45:931–933.
- Altemus M, Redwine LS, Leong Y-M, Frye CA, Porges SW, Carter CS (2001): Responses to laboratory psychosocial stress in postpartum women. *Psychosom Med* 63:814–821.
- Amico JA, Challinor SM, Cameron JL (1990): Pattern of oxytocin concentrations in the plasma and cerebrospinal fluid of lactating rhesus monkeys (*Macaca mulatta*): Evidence for functionally independent oxytocinergic pathways in primates. *J Clin Endocrinol Metab* 71:1531–1535.
- Amico JA, Johnston JM, Vagnucci AH (1994): Suckling-induced attenuation of plasma cortisol concentrations in postpartum lactating women. *Endocr Res* 20:79–87.
- Anderberg UM, Uvnas-Moberg K (2000): Plasma oxytocin levels in female fibromyalgia syndrome patients. *Z Rheumatol* 59:373–379.
- Born J, Lange T, Kern W, McGregor GP, Bickel U, Fehm HL (2002): Sniffing neuropeptides: A transnasal approach to the human brain. *Nat Neurosci* 5:514–516.
- Born J, Pietrowsky R, Fehm HL (1998): Neuropsychological effects of vasopressin in healthy humans. *Prog Brain Res* 119:619–643.
- Broadhead WE, Kaplan BH, James SA, Wagner EH, Schoenbach VJ, Grimson R, et al (1983): The epidemiologic evidence for a relationship between social support and health. *Am J Epidemiol* 117:521–537.
- Brownstein MJ, Russell JT, Gainer H (1980): Synthesis, transport, and release of posterior pituitary hormones. *Science* 207:373–378.
- Bruce ML (2002): Psychosocial risk factors for depressive disorders in late life. *Biol Psychiatry* 52:175–184.
- Bruins J, Hijman R, Van Ree JM (1992): Effect of a single dose of des-glycinamide-[Arg8]vasopressin or oxytocin on cognitive processes in young healthy subjects. *Peptides* 13:461–468.
- Buchanan J (1995): Social support and schizophrenia: A review of the literature. *Arch Psychiatr Nursing* 9:68–76.
- Carter CS (1998): Neuroendocrine perspectives on social attachment and love. *Psychoneuroendocrinology* 23:779–818.
- Carter CS, Altemus M (1997): Integrative functions of lactational hormones in social behavior and stress management. *Ann NY Acad Sci* 807:164–174.
- Chiodera P, Salvarani C, Bacchi-Modena A, Spallanzani R, Cigarini C, Alboni A, et al (1991): Relationship between plasma profiles of oxytocin and adrenocorticotrophic hormone during suckling or breast stimulation in women. *Horm Res* 35:119–123.
- Demitrack MA, Lesem MD, Listwak SJ, Brandt HA, Jimerson DC, Gold PW (1990): CSF oxytocin in anorexia nervosa and bulimia nervosa: Clinical and pathophysiological considerations. *Am J Psychiatry* 147:882–886.
- den Boer JA, Westenberg HG (1992): Oxytocin in obsessive compulsive disorder. *Peptides* 13:1083–1085.
- de Wied D, Diamant M, Fodor M (1993): Central nervous system effects of the neurohypophyseal hormones and related peptides. *Front Neuroendocrinol* 14:251–302.

- Domes G, Heinrichs M, Reichwald U, Hautzinger M (2002): Hypothalamic-pituitary-adrenal axis reactivity to psychological stress and memory in middle-aged women: High responders exhibit enhanced declarative memory performance. *Psychoneuroendocrinology* 27:843–853.
- Dressendorfer RA, Kirschbaum C, Rohde W, Stahl F, Strasburger CJ (1992): Synthesis of a cortisol-biotin conjugate and evaluation as tracer in an immunoassay for salivary cortisol measurement. *J Steroid Biochem Mol Biol* 43:683–692.
- Ehlert U, Straub R (1998): Physiological and emotional response to psychological stressors in psychiatric and psychosomatic disorders. *Ann N Y Acad Sci* 851:477–486.
- Engelmann M, Wotjak CT, Ebner K, Landgraf R (2000): Behavioural impact of intraseptally released vasopressin and oxytocin in rats. *Exp Physiol* 85(Spec No):125S–130S.
- Epperson CN, McDougle CJ, Price LH (1996): Intranasal oxytocin in obsessive-compulsive disorder. *Biol Psychiatry* 40:547–549.
- Erickson DH, Beiser M, Iacono WG (1998): Social support predicts 5-year outcome in first-episode schizophrenia. *J Abnorm Psychol* 107:681–685.
- Evans O, Steptoe A (2001): Social support at work, heart rate, and cortisol: A self-monitoring study. *J Occup Health Psychol* 6:361–370.
- Fehm-Wolfsdorf G, Bachholz G, Born J, Voigt K, Fehm HL (1988): Vasopressin but not oxytocin enhances cortical arousal: An integrative hypothesis on behavioral effects of neurohypophyseal hormones. *Psychopharmacology* 94:496–500.
- Frank GK, Kaye WH, Altemus M, Greeno CG (2000): CSF oxytocin and vasopressin levels after recovery from bulimia nervosa and anorexia nervosa, bulimic subtype. *Biol Psychiatry* 48:315–318.
- Gallo LC, Smith TW, Kircher JC (2000): Cardiovascular and electrodermal responses to support and provocation: Interpersonal methods in the study of psychophysiological reactivity. *Psychophysiology* 37:289–301.
- Gerin W, Pieper C, Levy R, Pickering TG (1992): Social support in social interaction: A moderator of cardiovascular reactivity. *Psychosom Med* 54:324–336.
- Green L, Fein D, Modahl C, Feinstein C, Waterhouse L, Morris M (2001): Oxytocin and autistic disorder: Alterations in peptide forms. *Biol Psychiatry* 50:609–613.
- Hays JC, Steffens DC, Flint EP, Bosworth HB, George LK (2001): Does social support buffer functional decline in elderly patients with unipolar depression? *Am J Psychiatry* 158:1850–1855.
- Heinrichs M (2000): *Oxytocin and Behavior: Psychobiological Effects of Oxytocin on Human Cognitive Performance and Stress Reactivity*. Göttingen, Germany: Cuvillier.
- Heinrichs M, Meinlschmidt G, Neumann I, Wagner S, Kirschbaum C, Ehlert U, Hellhammer DH (2001): Effects of suckling on hypothalamic-pituitary-adrenal axis responses to psychosocial stress in postpartum lactating women. *J Clin Endocrinol Metab* 86:4798–4804.
- Heinrichs M, Neumann I, Ehlert U (2002): Lactation and stress: Protective effects of breast-feeding in humans. *Stress* 5:195–203.
- Hellhammer DH, Buchtal J, Gutberlet I, Kirschbaum C (1997): Social hierarchy and adrenocortical stress reactivity in men. *Psychoneuroendocrinology* 22:643–650.
- Higuchi T, Negoro H, Arita J (1989): Reduced responses of prolactin and catecholamine to stress in the lactating rat. *J Endocrinol* 122:495–498.
- House JS, Landis KR, Umberson D (1988): Social relationships and health. *Sci* 241:540–545.
- Insel TR (1997): A neurobiological basis of social attachment. *Am J Psychiatry* 154:726–735.
- Insel TR, Young LJ (2001): The neurobiology of attachment. *Nat Rev Neurosci* 2:129–136.
- Kirschbaum C, Klauer T, Filipp SH, Hellhammer DH (1995): Sex-specific effects of social support on cortisol and subjective responses to acute psychological stress. *Psychosom Med* 57:23–31.
- Kirschbaum C, Kudielka BM, Gaab J, Schommer NC, Hellhammer DH (1999): Impact of gender, menstrual cycle phase, and oral contraceptives on the activity of the hypothalamus-pituitary-adrenal axis. *Psychosom Med* 61:154–162.
- Kirschbaum C, Pirke KM, Hellhammer DH (1993): The ‘Trier Social Stress Test’—a tool for investigating psychobiological stress responses in a laboratory setting. *Neuropsychobiology* 28:76–81.
- Knox SS, Uvnas-Moberg K (1998): Social isolation and cardiovascular disease: An atherosclerotic pathway? *Psychoneuroendocrinology* 23:877–890.
- Kraemer HC, Stice E, Kazdin A, Offord D, Kupfer D (2001): How do risk factors work together? Mediators, moderators, and independent, overlapping, and proxy risk factors. *Am J Psychiatry* 158:848–856.
- Laireiter AR (1996): *Skalen Soziale Unterstützung SSU [Social Support Scales Manual]*. Testmanual. Mödling, Germany: Dr. Schuhfried.
- Laux L, Glanzmann P, Schaffner P, Spielberger CD (1981): *Das State-Trait-Angstinventar. Theoretische Grundlagen und Handanweisungen [Manual for the State-Trait-Anxiety Inventory]*. Weinheim, Germany: Beltz.
- Leckman JF, Goodman WK, North WG, Chappell PB, Price LH, Pauls DL, et al (1994): Elevated cerebrospinal fluid levels of oxytocin in obsessive-compulsive disorder. Comparison with Tourette’s syndrome and healthy controls. *Arch Gen Psychiatry* 51:782–792.
- Legros JJ (2001): Inhibitory effect of oxytocin on corticotrope function in humans: Are vasopressin and oxytocin ying-yang neurohormones? *Psychoneuroendocrinology* 26:649–655.
- Lepore SJ, Allen KA, Evans GW (1993): Social support lowers cardiovascular reactivity to an acute stressor. *Psychosom Med* 55:518–524.
- Light KC, Smith TE, Johns JM, Brownley KA, Hofheimer JA, Amico JA (2000): Oxytocin responsivity in mothers of infants: A preliminary study of relationships with blood pressure during laboratory stress and normal ambulatory activity. *Health Psychol* 19:560–567.
- Lightman SL, Young WS (1989): Lactation inhibits stress-mediated secretion of corticosterone and oxytocin and hypothalamic accumulation of corticotropin-releasing factor and enkephalin messenger ribonucleic acids. *Endocrinology* 124:2358–2364.
- Macdonald EM, Jackson HJ, Hayes RL, Baglioni AJJ, Madden C (1998): Social skill as determinant of social networks and

- perceived social support in schizophrenia. *Schizophr Res* 29:275–286.
- MacMahon KM, Lip GY (2002): Psychological factors in heart failure: A review of the literature. *Arch Intern Med* 162:509–516.
- Martin A, State M, Anderson GM, Kaye WM, Hanchett JM, McConaha CW, et al (1998): Cerebrospinal fluid levels of oxytocin in Prader-Willi syndrome: A preliminary report. *Biol Psychiatry* 44:1349–1352.
- McCarthy MM, Altemus M (1997): Central nervous system actions of oxytocin and modulation of behavior in humans. *Mol Med Today* 3:269–275.
- Modahl C, Green L, Fein D, Morris M, Waterhouse L, Feinstein C, Levin H (1998): Plasma oxytocin levels in autistic children. *Biol Psychiatry* 43:270–277.
- Neumann I, Ludwig M, Engelmann M, Pittman QJ, Landgraf R (1993): Simultaneous microdialysis in blood and brain: Oxytocin and vasopressin release in response to central and peripheral osmotic stimulation and suckling in the rat. *Neuroendocrinology* 58:637–645.
- Neumann ID, Johnstone HA, Hatzinger M, Liebsch G, Shipston M, Russell JA, et al (1998): Attenuated neuroendocrine responses to emotional and physical stressors in pregnant rats involve adenohypophysial changes. *J Physiol* 508:289–300.
- Neumann ID, Krömer SA, Toschi N, Ebner K (2000a): Brain oxytocin inhibits the (re)activity of the hypothalamo-pituitary-adrenal axis in male rats: Involvement of hypothalamic and limbic brain regions. *Regul Pept* 96:31–38.
- Neumann ID, Torner L, Wigger A (2000b): Brain oxytocin: Differential inhibition of neuroendocrine stress responses and anxiety-related behaviour in virgin, pregnant and lactating rats. *Neuroscience* 95:567–575.
- Nissen E, Uvnas-Moberg K, Svensson K, Stock S, Widstrom AM, Winberg J (1996): Different patterns of oxytocin, prolactin but not cortisol release during breastfeeding in women delivered by caesarean section or by the vaginal route. *Early Hum Dev* 45:103–118.
- Paykel ES (2001): Stress and affective disorders in humans. *Semin Clin Neuropsychiatry* 6:4–11.
- Pedersen CA (1997): Oxytocin control of maternal behavior. Regulation by sex steroids and offspring stimuli. *Ann N Y Acad Sci* 807:126–145.
- Pedersen CA, Boccia ML (2002): Oxytocin links mothering received, mothering bestowed and adult stress responses. *Stress* 5:159–267.
- Pitman RK, Orr SP, Lasko NB (1993): Effects of intranasal vasopressin and oxytocin on physiologic responding during personal combat imagery in Vietnam veterans with posttraumatic stress disorder. *Psychiatry Res* 48:107–117.
- Rozanski A, Blumenthal JA, Kaplan J (1999): Impact of psychological factors on the pathogenesis of cardiovascular disease and implications for therapy. *Circulation* 99:2192–2217.
- Ruberman W, Weinblatt E, Goldberg JD, Chaudhary BS (1984): Psychosocial influences on mortality after myocardial infarction. *N Engl J Med* 311:552–559.
- Russell JA, Douglas AJ, Ingram CD (2001): Brain preparations for maternity—adaptive changes in behavioral and neuroendocrine systems during pregnancy and lactation. An overview. *Prog Brain Res* 133:1–38.
- Sayal K, Checkley S, Rees M, Jacobs C, Harris T, Papadopoulos A, Poon L (2002): Effects of social support during weekend leave on cortisol and depression ratings: A pilot study. *J Affect Disord* 71:153–157.
- Schwarzer R, Leppin A (1991): Social support and health: A theoretical and empirical overview. *J Personal Soc Relationships* 8:99–127.
- Seeman TE (2000): Health promoting effects of friends and family on health outcomes in older adults. *Am J Health Promotion* 14:362–370.
- Seeman TE, McEwen BS (1996): Impact of social environment characteristics on neuroendocrine regulation. *Psychosom Med* 58:459–471.
- Smith TW, Ruiz JM (2002): Psychosocial influences on the development and course of coronary heart disease: Current status and implications for research and practice. *J Consult Clin Psychol* 70:548–568.
- Spitzer SB, Llabre MM, Ironson GH, Gellman MD, Schneiderman N (1992): The influence of social situations on ambulatory blood pressure. *Psychosom Med* 54:79–86.
- Stephens A (2000): Stress, social support and cardiovascular activity over the working day. *Int J Psychophysiol* 37:299–308.
- Steyer R, Schwenkmezger P, Notz P, Eid M (1997): *Der Mehrdimensionale Befindlichkeitsfragebogen (MDBF)* [Multidimensional Mood Questionnaire]. Göttingen, Germany: Hogrefe.
- Swaab DF, Pool CW, Nijveldt F (1975): Immunofluorescence of vasopressin and oxytocin in the rat hypothalamo-neurohypophysial system. *J Neural Transm* 36:195–215.
- Turner RA, Altemus M, Enos T, Cooper B, McGuinness T (1999): Preliminary research on plasma oxytocin in normal cycling women: Investigating emotion and interpersonal distress. *Psychiatry* 62:97–113.
- Uchino BN, Cacioppo JT, Kiecolt-Glaser JK (1996): The relationship between social support and physiological processes: A review with emphasis on underlying mechanisms and implications for health. *Psychol Bull* 119:488–531.
- Uchino BN, Garvey TS (1997): The availability of social support reduces cardiovascular reactivity to acute psychological stress. *J Behav Med* 20:15–27.
- Uchino BN, Holt-Lunstad J, Uno D, Betancourt R, Garvey TS (1999): Social support and age-related differences in cardiovascular function: An examination of potential mediators. *Ann Behav Med* 21:135–142.
- Uno D, Uchino BN, Smith TW (2002): Relationship quality moderates the effect of social support given by close friends on cardiovascular reactivity in women. *Int J Behav Med* 9:243–262.
- Uvnas-Moberg K (1998): Oxytocin may mediate the benefits of positive social interaction and emotions. *Psychoneuroendocrinology* 23:819–835.
- Uvnas-Moberg K, Ahlenius S, Hillegaart V, Alster P (1994): High doses of oxytocin cause sedation and low doses cause and anxiolytic-like effect in male rats. *Pharmacol Biochem Behav* 49:101–106.
- Uvnas-Moberg K, Björkstrand E, Hillegaart V, Ahlenius S (1999): Oxytocin as a possible mediator of SSRI-induced antidepressant effects. *Psychopharmacology* 142:95–101.

- Vandesande F, Dierickx K (1975): Identification of the vasopressin producing and of the oxytocin producing neurons in the hypothalamic magnocellular neurosecretory system of the rat. *Cell Tissue Res* 164:153–162.
- Veiel HOF, Baumann U (1992): *The Meaning and Measurement of Social Support*. Washington: Hemisphere.
- Vining RF, McGinley RA (1987): The measurement of hormones in saliva: Possibilities and pitfalls. *J Steroid Biochem* 27:81–94.
- Walker CD, Lightman SL, Steele MK, Dallman MF (1992): Suckling is a persistent stimulus to the adrenocortical system of the rat. *Endocrinology* 130:115–125.
- Walker CD, Trottier G, Rochford J, Lavalley D (1995): Dissociation between behavioral and hormonal responses to the forced swim stress in lactating rats. *J Neuroendocrinol* 7:615–622.
- Windle RJ, Shanks N, Lightman SL, Ingram CD (1997a): Central oxytocin administration reduces stress-induced corticosterone release and anxiety behavior in rats. *Endocrinology* 138:2829–2834.
- Windle RJ, Wood S, Shanks N, Perks P, Conde GL, da Costa AP, et al (1997b): Endocrine and behavioural responses to noise stress: Comparison of virgin and lactating female rats during non-disrupted maternal activity. *J Neuroendocrinol* 9:407–414.
- Winer BJ (1971): *Statistical Principles in Experimental Design*, 2nd ed. New York: McGraw-Hill.
- Wotjak CT, Ganster J, Kohl G, Holsboer F, Landgraf R, Engelmann M (1998): Dissociated central and peripheral release of vasopressin, but not oxytocin, in response to repeated swim stress: New insights into the secretory capacities of peptidergic neurons. *Neuroscience* 85:1209–1222.
- Young EA, Lopez JF, Murphy-Weinberg V, Watson SJ, Akil H (2000): Hormonal evidence for altered responsiveness to social stress in major depression. *Neuropsychopharmacology* 23:411–418.
- Zung WWK (1965): A self-rating depression scale. *Arch Gen Psychiatry* 12:63–70.